

Long Beach, California — 1999 (N=83)

Figure A. Age of GISP participants, in years, 1999




Figure B. Race/ethnicity of GISP participants, 1999




Figure C. Percentage of GISP participants identifying as men who have sex with men, 1988–1999





Figure D. Reason for visit among GISP participants, 1999



Long Beach, California — 1999 (N=83)

Figure E. Previous episode of gonorrhea among GISP participants, 1991–1999




Figure F. Drugs used to treat gonorrhea among GISP participants, 1999




Figure G. Drugs used to treat chlamydia trachomatis infection among GISP participants, 1999





Figure H. Resistance to penicillin and tetracycline among GISP isolates, 1999



Long Beach, California — 1999 (N=83)

Figure I. Decreased susceptibility to ceftriaxone among GISP isolates, 1988–1999

No isolates with decreased susceptibility to ceftriaxone have been identified at this clinic.

Figure J. Decreased susceptibility to cefixime among GISP isolates, 1992–1999






Figure K. Intermediate resistance and resistance to ciprofloxacin among GISP isolates, 1990–1999



Note: Susceptibility to ciprofloxacin first measured in 1990.

Figure L. Decreased susceptibility to azithromycin among GISP isolates, 1992–1999



Note: Susceptibility to azithromycin first measured in 1992.

Note: Decreased susceptibility to azithromycin is defined here as $>$ or $=$ 1.0 ug/ml.
No NCCLS criteria currently exist.